Patient Information

How to do REVERSE PRESSURE SOFTENING?

Developed by K. Jean Cotterman RNC-E, IBCLC

Try this in the early weeks of learning to breastfeed, if pain, swelling, or fullness cause problems:

The <u>key</u> is making the areola (the circle) <u>very soft</u> right around the base of the nipple, <u>right before every feeding</u>, for better latching.

- Your baby needs the areola soft to get a "good mouthful" of breast, beyond the nipple, to <u>remove more milk</u>, and be very gentle to your nipple.
- Lie most of the way down on your back when you soften the areola. (This is a great way for your baby and you to learn to breastfeed too).
- Look at the pictures below. Choose the easiest way for you to press on the areola.
- Be gentle to avoid pain, but press the circle firmly and steadily in toward your heart, <u>counting slowly to 50;</u> *Count very slowly if very swollen*.
- In the hospital, ask someone to show you. At home, ask for help from a relative or friend.
- Soften the areola right before each feeding (or pumping) till latching is always easy.

If you need to pump milk for your baby,

- Always soften the areola first before you pump, too.
- Use only medium or low vacuum so swelling won't move back into the areola.
- Pause to re-soften the areola once or twice, to get more **total** milk out.
- Make pumping sessions short, but often, (2-3 hours.) while awake.

<u>Method 1: Two step method</u>, use the straight fingers of both hands, placed with the knuckles touching the nipple: Count to 50 in each position. **If very swollen, count very slowly.**



All drawings by Kyle Cotterman



1. Place fingers on each side of the nipple.

2. Place fingers above and below the nipple

Method 2: Soft Ring Method: cut off bottom half of an artificial nipple to place on the areola and press with fingers. Count to 50. If very swollen, count very slowly.





Method 3: One handed "flower hold". Fingernails, short, fingertips curved, placed where baby's tongue will go. Count to 50. If very swollen, count very slowly.



Distributed by the Middlesex London Breastfeeding Coalition, January 2014. Currently reviewed by the Middlesex London Elgin Breastfeeding Coalition with with representation from London Health Sciences Centre, Strathroy Middlesex General Hospital Southwestern Public Health-St. Thomas, St.

Thomas Elgin general Hospital and the Middlesex London Health Unit, November 2018.

This handout may be copied and distributed without further permission, on the condition that it is not used in <u>any</u> context violating

The International WHO Code on the Marketing of Breastmilk Substitutes